



Tradecorp AZ Fresco

**100% soluble micro-granule to
prevent & correct multiple
deficiencies**

Description

Premium 6 micro-nutrient EDTA chemical mix with emphasis on Iron (Fe) 4%, Zinc (Zn) 4% and Manganese (Mn) 3%

- Corrects and prevents micro-nutrient deficiencies in all crops
- EDTA Chelates with maximum Chelation %*
- 100% soluble micro-granule (WG) for rapid & instantaneous solubility
- Suitable for use in foliar, drip, direct soil, substrate & hydroponic applications
- Suitable for use in soils & solutions with pH 4-7
- Production system certified to ISO 9001 & ISO 14001
- Guaranteed Contents, Ecocert Approved
- 5 kg bag size
- (* Chelation % is the % of the micronutrient attached to the EDTA, higher is better)

Benefits

- Chemical premix that tackles all micronutrient deficiencies
- EDTA Chelates are the most efficient form of nutrient for soil and foliar use
- EDTA Chelates remain soluble in the soil solution for continued availability
- Compatible with most commonly used fertilizers and agrochemical products
- Simple and convenient handling without lumps or sediments
- Easy pouring micro-granule (WG), no rinsing of packing
- High efficiency at low dose rates

Technical Specifications

- Magnesium (Mg) soluble in water: 1.20% w/w
- Sulfur (S) soluble in water: 0.96% w/w
- Zinc (Zn) soluble in water and chelated by EDTA: 4.0% w/w
- Iron (Fe) soluble in water and chelated by EDTA: 4.0% w/w
- Manganese (Mn) soluble in water and chelated by EDTA: 3.0% w/w
- Copper (Cu) soluble in water and chelated by EDTA: 0.60% w/w
- Boron (B) soluble in water: 1.50% w/w
- Molybdenum (Mo) soluble in water: 0.05% w/w
- EDTA pH stability range: 4-7

Rovensa ANZ

PO BOX 2046
Graceville QLD
1300 595 000

Info.au@rovensanext.com
www.rovensanext-au-nz.com



Let's grow greener



Rovensanext

Tradecorp AZ Fresco

Dosage and method of use

Foliar Application		
General Dose	0.5 – 1.5 Kg/ Ha	2-3 applications before or as soon as symptoms appear 14 days apart
Vegetables (e.g. tomato, cucurbits)	0.5 – 1 Kg/ Ha (75 – 100g / 100L)	2-3 applications early in the crop cycle once foliage is sufficient
Tubers (Potato)	1 Kg/ Ha	3-4 applications once stems reach 10 cm. Final application post flowering. Use high volumes of water (e.g. 500 L)
Orchard Crops	1 – 1.5 Kg /Ha	2-4 applications before flowering x2-3 and during fruit set x1
Cereals	0.5 – 1 Kg/ Ha	2-3 applications at start of tillering, stem elongation and flag leaf
Ornamentals	0.2 – 0.5 Kg/ Ha (75 – 100 g/ 100L)	2-3 applications every 7-10 days from beginning of vegetative cycle
Other Applications		
Fertigation - General Dose	2 – 3 Kg/ Ha	2-4 applications every 7-10 days from early in the crop cycle
Hydroponics - General Dose	3 – 5 Kg/ 1000L (x 100)	Use 1 L of mother stock solution /100 L irrigation water. Use the higher rate from fruit set, high temperatures or shorter winter days
Soil Applications - General Dose	5 – 10 Kg/ Ha	1 application before sowing, transplant or the beginning of plant growth